


AD-A193 383 MOLECULAR BIOLOGY OF ARCHAEABACTERIA(U) BRITISH COLUMBIA 1/1  
UNIV VANCOUVER DEPT OF BIOCHEMISTRY P P DENNIS  
31 MAR 88 N00014-87-G-0049

UNCLASSIFIED

F/G 6/2

NL





AD-A193 383

## REPORT DOCUMENTATION PAGE

DTIC FILE COPY

②

1a. RESTRICTIVE MARKINGS			1b. RESTRICTIVE MARKINGS		
2a. DECLASSIFICATION / DOWNGRADING SCHEDULE			3. DISTRIBUTION / AVAILABILITY OF REPORT		
NA			DISTRIBUTION STATEMENT A		
2b. DECLASSIFICATION / DOWNGRADING SCHEDULE			Approved for public release Distribution Unlimited		
4. PERFORMING ORGANIZATION REPORT NUMBER(S)			5. MONITORING ORGANIZATION REPORT NUMBER(S)		
NA					
6a. NAME OF PERFORMING ORGANIZATION		6b. OFFICE SYMBOL (If applicable)		7a. NAME OF MONITORING ORGANIZATION	
Univ. of British Columbia				DTIC ELECTE	
6c. ADDRESS (City, State, and ZIP Code)		7b. ADDRESS (City, State, and ZIP Code)		MAR 29 1988	
Department of Biochemistry 2146 Health Sciences Mall Vancouver, B.C. V6T 1W5				S D	
8a. NAME OF FUNDING / SPONSORING ORGANIZATION		8b. OFFICE SYMBOL (If applicable)		9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER	
Office of Naval Research		ONR		N00014-87-G-0049	
8c. ADDRESS (City, State, and ZIP Code)		10. SOURCE OF FUNDING NUMBERS			
800 N. Quincy St. Arlington, VA 22217-5000		PROGRAM ELEMENT NO. 61153N		PROJECT NO. RR04106	
		TASK NO.		WORK UNIT ACCESSION NO.	
11. TITLE (Include Security Classification)					
Molecular Biology of Archaeobacteria					
12. PERSONAL AUTHOR(S)					
Patrick P. Dennis					
13a. TYPE OF REPORT		13b. TIME COVERED		14. DATE OF REPORT (Year, Month, Day)	
Annual		FROM 4/87 TO 3/88		880331	
15. PAGE COUNT					
16. SUPPLEMENTARY NOTATION					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB-GROUP			
19. ABSTRACT (Continue on reverse if necessary and identify by block number)					
<p>Genomic restriction fragments from <u>Halobacterium cutirubrum</u> (Hcu) and <u>Sulfolobus acidocaldarius</u> (Sac) encoding ribosomal protein L11e, L1e, L10e and L12e were cloned and sequenced. The sequences were analyzed and the reading frames identified. The deduced L10e and L12e protein sequences were aligned with eubacterial and eucaryotic proteins and their structure-function and evolutionary relationships were defined. The N-terminal amino acid sequence of the purified superoxide dismutase enzymes from Hcu was determined, an oligonucleotide was prepared and the gene was cloned and sequenced. The sequence has been aligned to other SOD sequences. The gene transcripts have been characterized and the regulation of transcription (induction) by oxygen radicals has been demonstrated and characterized.</p>					
20. DISTRIBUTION STATEMENT OF ABSTRACT			21. ABSTRACT SECURITY CLASSIFICATION		
<input type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS					
22a. NAME OF RESPONSIBLE INDIVIDUAL			22b. TELEPHONE (Include Area Code)		22c. OFFICE SYMBOL

Abbreviations: Hcu - Halobacterium cutirubrum  
 Hha - Halobacterium holobium (same species as Hcu)  
 Hvo - Halobacterium volcanii  
 Sac - Sulfolobus acidocaldarius  
 Eco - Escherichia coli

### Research Objectives

(i) to characterize the principles of gene organization and regulation of gene expression in archaeobacteria; (ii) to elucidate the evolutionary relationship between these novel organisms and the traditional eubacterial and eucaryotic organisms; (iii) to understand in biophysical and molecular terms some of the mechanisms that allow archaeobacteria to inhabit extreme environments.

### Progress - Year 1

Ribosomal protein genes: The ribosomal A protein complex forms the stalk structure on the large ribosome subunit and is comprised of four copies (two dimers) of L12e and one copy of L10e ribosomal protein. The L10e protein either binds to ribosomal RNA either directly or through the L11e protein which forms a bulge at the base of the stalk. This domain on the large subunit is the site of factor binding and associated GTPase activities during the protein synthesis cycle and is a conserved and defined feature on the ribosome from all organisms (1). In Eco the genes encoding the L11, L1, L10 and L12 ribosomal proteins are located within a 3.0 Kb region of genomic DNA. This region has been cloned, sequenced and extensively characterized (2,3).

Partial amino acid sequences of many archaeobacterial ribosomal proteins have been determined (4). We used the sequences from the Hcu L11e and L12e proteins to deduce and synthesize oligonucleotide sequences complementary to the corresponding genes. We cloned initially a 1.2 Kb Pst-Bam fragment encoding L12e and subsequently a 5.2 Kb Cla-Bam fragment containing both L11e and L12e. The sequence of the entire 5.2 Kb fragment was determined and it was found to contain several long open reading frames. Four of these open reading frames were shown to encode the L11e, L1e, L10e and L12e ribosomal proteins. The gene order as present in Eco is conserved in Hcu although the transcriptional organization will certainly be much different (work in progress).

Using a similar strategy the corresponding genomic region of Sac has been cloned and sequenced. It contains the same four genes in the same order. From the Hcu and Sac nucleic acid sequence we have deduced the amino acid sequences for the four ribosomal proteins and have begun to align the protein to their eubacterial and eucaryotic equivalent and deduce their evolutionary relationships. This analysis has been completed for the L10e and L12e protein (5).

There are presently four complete sequences of L10e proteins (one eubacterial, two archaeobacterial and one eucaryotic) and 16 complete sequences of L12e (eight eubacterial, three archaeobacterial and five eucaryotic). Using these sequences we have carried out inter-kingdom L10e and L12e alignments. The L10e protein from the three kingdoms were found to be co-linear. (A cDNA sequence published by Rich and Steitz was shown by us to encode the human L10e

protein (6)). The eubacterial protein is much shorter than the archaeobacterial and eucaryotic proteins because of two large deletions, one internal and one at the carboxy terminus. The two archaeobacterial L10e proteins were most similar; they exhibit 27% amino acid identity with only a single deletion-insertion. Inter-kingdom comparisons exhibit 15-25% amino acid identity with 5-7 deletion insertions.

The L12e protein from archaeobacteria and eucaryotes can also be aligned and are co-linear. The eubacterial protein could not be made to fit this pattern although two regions appear to have homologous domain in the archaeobacterial and eucaryotic proteins. The C-terminal domain of the Eco L12e protein has been shown to contain (i) a conserved face for interaction with extrinsic translation factors; (ii) an anion (potential GTP) binding site; and (iii) a putative L12e-L12e dimerization site (1). This domain is separated from the N-terminus by an unstructured ala-pro-rich hinge sequence and aligns to a region near the amino terminus of the archaeobacterial and eucaryotic proteins. Inter-kingdom comparisons range from 18-36 percent amino acid sequence identity over their domain. The second region conserved in all proteins is the ala-pro-rich sequence. In eubacteria this sequence precedes the factor interaction domain; in archaeobacteria and eucaryotic it follows the factor interaction domain. The N-terminus of the eubacterial L12e protein exhibits more similarity to its own carboxy end than to any region within the archaeobacterial-eucaryotic protein. The carboxy termini of the archaeobacterial and eucaryotic protein are highly similar to each other and apparently not represented in the eubacterial L12e protein.

Intraspecies comparisons between L10e and L12e proteins indicate that the L10e protein of archaeobacteria and eucaryotes contains a partial copy of the L12e protein fused to its carboxy terminus. (In eubacteria most of this fusion has been removed from L10e by the carboxy terminal deletion and the L12e protein has been restructured.) For the two Sac proteins the 31 carboxy terminal residues of L10e and L12e are identical; conservation at the nucleotide level is also perfect. This homology extends further into the central regions of the L10e and L12e proteins and includes a 26 amino acid long module, reiterated thrice in archaeobacterial L10e, twice in eucaryotic L10e and once in the corresponding L12e protein. Careful examination indicates that a single copy of the modular sequence is also present in the L12e and L10e protein of eubacteria. This modular sequence may play a role in L12e dimerization, L10e-L12e complex formation and the function of the L10e-L2e complex in translation.

Based upon these alignments and shared homologies a model has been constructed to depict the evolution of the primordial L10e and L12e genes and proteins in the three phylogenetic groups (5).

**Superoxide dismutase:** Previously we had purified and characterized the SOD enzyme activity from Hcu (7). An N-terminal amino acid sequence was determined, an oligonucleotide was synthesized and the gene encoding the SOD enzyme was cloned as a 1.1 Kb Sau3A fragment and sequenced. The fragment contain a single long open reading frame encoding a protein of 200 amino acids. The first 56 codons of the ORF correspond to the amino acid sequence determined from the purified protein. The gene encodes a typical Mn/Fe ion type enzyme as evidenced by the fact that (i) it has an acidic pI; (ii) it



pr HP

codes

and/or  
label

A-1

contains no cys residues; (iii) it is relatively abundant in tyr and trp residues; (iv) it exhibits 41% amino acid identity with the Mn SOD of *Bacillus stearothermophilus* and (v) it conserves the four residues, his 29, his 76, asp 158 and his 162 that in the *Bacillus* enzyme (position 26, 81, 163 and 167 respectively) are utilized for Mn binding.

By S1 nuclease protection and primer extension analysis we have shown that the SOD mRNA is about 242 nucleotides in length. It begins two nucleotides in front of the ATG translation initiation codon and terminates about 40 nucleotides downstream of the TAA translation termination codon, in a T<sub>5</sub> sequence that is preceded by a GC-rich sequence. This terminator sequence appears similar to other *Hcu* terminators. Analysis of this and other 5' flanking sequences have yet to reveal the consensus sequence involved in promoter recognition by RNA polymerase in *Hcu* (that is, there is little conservation between the bacterio-opsin, SOD, rRNA and r-protein 5' flanking sequences).

Like bacterio-opsin, the mRNA encoding SOD starts only two nucleotides in front of the ATG translation initiation codon. There is no homolog of the Shine Dalgarno purine-rich sequence of eubacteria that is complementary to the 3' end of 16S rRNA on the SOD mRNA. In contrast some r-protein genes of *Hcu* retain this sequence. Thus it seems that *Hcu* ribosomes can initiate translation by two separate mechanisms: at the first AUG on the transcript (SOD and bacterio-opsin) or by a Shine Dalgarno interaction at internal AUG codons.

We have discovered that the SOD gene in *Hcu* is subject to regulation. Addition of paraquat, a generator of oxygen radicals, to an exponential phase culture results in the 5-10x induction of SOD enzyme activity. After several days of growth in the presence of paraquat the SOD activity returns to normal. The culture has become resistant to the inhibitor and the frequency suggests that resistance arises by transposition insertion (as in bacterio-opsin negative mutants of *Hha*; 8) into a gene specifying sensitivity (i.e. transport of paraquat across the cell envelope). This observation is being investigated.

### References

- (1) Liljas, A. (1982) *Prog. Biophys. Mol. Biol.* 40, 161-228.
- (2) Post, L., Strycharz, G., Nomura, M., Lewis, H. and Dennis, P. (1979) *PNAS* 76, 1697-1701.
- (3) Downing, W. and Dennis, P. (1987) *J. Mol. Biol.* 194, 609-620.
- (4) Matheson, A. (1985) *The Bacteria* 8, 761-777.
- (5) Shimmin, L., Ramirez, C., Matheson, A. and Dennis, P. (1988) submitted for publication.
- (6) Rich, B. and Steitz, J. (1988) *Mol. Cell Biol.*, in press.
- (7) May, B. and Dennis, P. (1987) *J. Bacteriol.* 169, 1417-1422.
- (8) Betlach, M., Leong, D. and Boyer, H. (1986) *System. Appl. Microbiol.* 7, 83-89.

## Objectives for Year 2

### Gene encoding r-protein

- 1) Carry out inter-kingdom alignment and analysis of L1le and L1e protein;
- 2) Define the transcriptional organization of the L1le, L1e, L10e and L12e gene clusters in Hcu and Sac, map 5' and 3' transcript ends, cap 5' transcript ends and analyze for conserved promoter and terminator sequences.

### Superoxide dismutase

- 3) Cap mRNA and characterize induction by paraquat at the level of transcription;
- 4) Look for DNA binding protein to the SOD promoter region by in vivo foot-printing;
- 5) Clone and characterize SOD gene from a genetically stable strain Hvo, isolate paraquat-sensitive (null) and resistant (constitutive) mutants;
- 6) Utilize the transformation system of Hvo to characterize the genetic regulation of the SOD gene.

### Processing of rRNA

- 7) Characterize promoter and processing signals in the separate 16S-23S and 5S transcription units of Sac;
- 8) Utilize cloned copies of the 16S inverted repeat processing signal from Hcu to develop an assay for RNaseIII-like activity;
- 9) Attempt to purify Hcu RNaseIII-like activity, clone the gene and study substrate specificity by in vitro site-specific mutagenesis of cloned 16S inverted repeat sequence.

### Publications

May, B. and P. Dennis (1987) Superoxide dismutase from the extremely halophilic archaebacterium Halobacterium cutirubrum. J. Bacteriol. 169, 1417-1422.

Shimmin, L., C. Ramirez, A. Matheson and P. Dennis (1988) Sequence alignment and evolutionary comparison of the L10e and L2e ribosomal proteins from archaebacteria, eubacteria and eucaryotics. Submitted for publication (copy enclosed to ONR).

### Training

Bruce May: MRC, predoctoral fellow, Canadian  
 Lawrence Shimmin: ONR, predoctoral student, Canadian  
 Craig Newton: ONR technician, Canadian  
 Diedre de Jong-Wong: ONR technician, Canadian  
 Phelgun Joshi: MRC predoctoral student (grant), Canadian  
 Peter Durovic: MRC predoctoral student (grant), Canadian

MOLECULAR BIOLOGY PROGRAM DISTRIBUTION L

ANNUAL, FINAL, AND TECHNICAL REPORTS (On

Dr. C. E. Ballou  
Department of Biochemistry  
University of California  
Berkeley, CA 94720

Dr. Rita Colwell  
Department of Microbiology  
University of Maryland  
College Park, MD 20742

Dr. Lacy Daniels  
Department of Microbiology  
University of Iowa  
Iowa City, IA 52242

Dr. Patrick P. Dennis  
Department of Biochemistry  
University of British Columbia  
2146 Health Sciences Mall  
Vancouver, B.C. V6T 1W5

Dr. Henryk Eisenberg  
The Weizmann Institute of Science  
Department of Polymer Research  
P.O. Box 26  
Rehovot 76100, Israel

Dr. James G. Ferry  
Department of Anaerobic Microbiology  
Virginia Polytechnic Institute  
and State University  
Blacksburg, Virginia 24061

Dr. Robert P. Gunsalus  
Department of Microbiology  
UCLA  
405 Hilgard Avenue  
Los Angeles, CA 90024

Dr. Ramesh Gupta  
Southern Illinois University  
Department of Chemistry and Biochemistry  
Carbondale, IL 62901

Dr. Holger W. Jannasch  
Woods Hole Oceanographic Institution  
Woods Hole, MA 02543

Dr. Jordan Konisky  
University of Illinois  
809 South Wright Street  
Champaign, IL 61820

Dr. Mary E. Lidstrom  
Center for Great Lakes Studies  
University of Wisconsin-Milwaukee  
Milwaukee, WI 53201

Dr. Ralph Mitchell  
Division of Applied Sciences  
Harvard University  
125 Pierce Hall  
Cambridge, MA 02138

Dr. Leonard Muscatine  
Department of Biology  
University of California  
Los Angeles, California 90024

Dr. David P. Nagle  
Department of Botany and Microbiology  
University of Oklahoma  
Norman, OK 73019

Dr. Gary J. Olsen  
Department of Biology  
Indiana University  
Jordan Hall 138  
Bloomington, Indiana 47405

Dr. Gregory J. Olson  
Inorganic Materials Division  
Center for Materials Science  
National Bureau of Standards  
Washington, DC 20234

Dr. Leo Parks  
Department of Microbiology  
North Carolina State University  
Raleigh, NC 27695

Dr. John N. Reeve  
Department of Microbiology  
Ohio State University  
484 West 12th Avenue  
Columbia, OH 43210-1292

Dr. V. Romanovsky  
Office d'Etudes Marines et Atmospherique  
64 rue Gabrielle  
1180, Bruxelles, Belgium

Dr. Saul Roseman  
Department of Biology  
Johns Hopkins University  
Baltimore, MD 21218

Dr. David C. White  
Department of Microbiology  
Institute of Applied Microbiology  
University of Tennessee  
Knoxville, TN 37996-0845

Dr. Carl R. Woese  
Genetics Department  
University of Illinois  
505 S. Goodwin Avenue  
Urbana, IL 61801

Dr. Ralph S. Wolfe  
131 Burrill Hall  
University of Illinois  
Urbana, IL 61801

Dr. Eli D. Schmel, Code 1141MB  
Office of Naval Research  
800 North Quincy Street  
Arlington, VA 22217-5000

Dr. Michael T. Marron, Code 1141MB  
Office of Naval Research  
800 North Quincy Street  
Arlington, VA 22217-5000

Dr. Margo G. Haygood  
Office of Naval Research  
800 North Quincy Street  
Arlington, VA 22217-5000

Administrator (2 copies, Enclose DTIC Form 50)  
Defense Technical Information Center  
Building 5, Cameron Station  
Alexandria, VA 22314

ANNUAL AND FINAL REPORTS ONLY (One copy each)

Commander  
Chemical and Biological Sciences Division  
Army Research Office  
P. O. Box 12211  
Research Triangle Park, NC 27709



Directorate of Life Sciences  
Air Force Office of Scientific Research  
Bolling Air Force Base  
Washington, DC 20332

Chemistry and Atmospheric Sciences Directorate  
Air Force Office of Scientific Research  
Bolling Air Force Base  
Washington, DC 20332

Director  
Biotechnology Division  
CRDEC  
Aberdeen Proving Grounds, MD 21010

Administrative Contracting Officer  
ONR Resident Representative  
(Address varies - obtain from your business office)

Director, Code 12  
Applied Research and Technology Directorate  
Office of Naval Research  
800 North Quincy Street  
Arlington, VA 22217-5000

Director, Code 22  
Support Technology Directorate  
Office of Naval Technology  
800 North Quincy Street  
Arlington, VA 22217-5000

Director, Code 112  
Environmental Sciences Directorate  
Office of Naval Research  
800 North Quincy Street  
Arlington, VA 22217-5000

Director, Code 113  
Chemistry Division  
Office of Naval Research  
800 North Quincy Street  
Arlington, VA 22217-5000

FINAL AND TECHNICAL REPORTS ONLY  
Director (6 copies)  
Naval Research Laboratory  
Attn: Technical Information Division, Code 2627  
Washington, DC 20375

END

DATE

FILMED

7-88

Dtic